AMENDMENTS TO THE CLAIMS:

Set forth below in ascending order, with status identifiers, is a complete listing of all claims currently under examination. Changes to any amended claims are indicated by strikethrough and underlining. This listing also reflects any cancellation and/or addition of claims.

1 to 52. (Canceled)

53. (Currently Amended) A method of producing a multi-layered tissue construct *in vivo* comprising the steps of:

providing a first polymerizable mixture which comprises first cells;

providing a second polymerizable mixture;

placing a volume of the first polymerizable mixture in an osteochondral tissue space wherein the first polymerizable mixture is in direct contact with the osteochondral tissue underneath, then crosslinking the first polymerizable mixture until the first polymerizable mixture forms an at least partially gelled first layer; ~~and~~

placing a volume of the second polymerizable mixture in the tissue space with the at least partially gelled first layer, then crosslinking the second polymerizable mixture until the second polymerizable mixture is at least partially gelled to form a second layer; and

additionally crosslinking the first and second polymerizable mixtures until the first and second layers further polymerize to form an integrated multi-layered gel.

54. (Previously Presented) The method according to claim 53, wherein the first polymerizable mixture comprises a polymerization initiator.

55. (Previously Presented) The method according to claim 53, wherein the second polymerizable mixture comprises a polymerization initiator.

56. (Previously Presented) The method according to claim 53, wherein the second polymerizable mixture comprises second cells.

57. (Previously Presented) The method according to claim 56, wherein the first and second cells are different cell types.

58. (Previously Presented) The method according to claim 57, wherein the first cells are mesenchymal stem cells and the second cells are chondrocytes.

59. (Previously Presented) The method according to claim 56, wherein the first and second cells are same cell types.

60. (Previously Presented) The method according to claim 59, wherein the first and second cells are chondrocytes.

61. (Previously Presented) The method according to claim 53, wherein the cells are autologous cells.

62. (Previously Presented) The method according to claim 53, further comprising the step of adding a suspension of cells to a surface of the at least partially gelled first layer, before the step of placing the volume of the second polymerizable mixture in the tissue space.

63. (Previously Presented) The method according to claim 53, wherein at least one of the first and second polymerizable mixtures further comprises a bioactive substance selected from the group consisting of a nutrient, a cellular mediator, a growth factor, a compound which induces cellular differentiation, a bioactive polymer, a gene vector, and a pharmaceutical.

64. (Canceled)

65. (Currently Amended) The method according to claim 53, wherein crosslinking of the first polymerizable mixture is controlled by exposure to an external radiation, and crosslinking of the second polymerizable mixture is controlled by exposure to the external radiation.

66. (Currently Amended) A method of producing a multi-layered tissue construct *in vivo* comprising the steps of:

providing a first polymerizable mixture which comprises first cells;

providing a second polymerizable mixture;

providing a third polymerizable mixture;

wherein at least one of the second and third polymerizable mixtures comprises cells;

placing a volume of the first polymerizable mixture in an osteochondral tissue space wherein the first polymerizable mixture is in direct contact with the osteochondral tissue underneath, then crosslinking the first polymerizable mixture until the first polymerizable mixture forms an at least partially gelled first layer;

placing a volume of the second polymerizable mixture in the tissue space with the at least partially gelled first layer, then crosslinking the second polymerizable mixture until the second polymerizable mixture is at least partially gelled to form a second layer;

placing a volume of the third polymerizable mixture in the tissue space with the at least partially gelled first and second layers, then crosslinking the third polymerizable mixture until the third polymerizable mixture is at least partially gelled to form a third layer; and

additionally crosslinking the first, second, and third polymerizable mixtures until the first, second, and third layers further polymerize to form an integrated multi-layered gel.

67. (Previously Presented) The method according to claim 66, wherein the first polymerizable mixture comprises a polymerization initiator.

68. (Previously Presented) The method according to claim 66, wherein the second polymerizable mixture comprises a polymerization initiator.

69. (Previously Presented) The method according to claim 66, wherein the third polymerizable mixture comprises a polymerization initiator.

70. (Previously Presented) The method according to claim 66, wherein the second polymerizable mixture comprises second cells, and the third polymerizable mixture comprises third cells.

71. (Previously Presented) The method according to claim 70, wherein the first, second, and third cells are different cell types.

72. (Previously Presented) The method according to claim 71, wherein the first cells are mesenchymal stem cells and the second cells are chondrocytes.

73. (Previously Presented) The method according to claim 70, wherein the first, second, and third cells are same cell types.

74. (Previously Presented) The method according to claim 73, wherein the first, second, and third cells are chondrocytes.

75. (Previously Presented) The method according to claim 66, wherein the cells are autologous cells.

76. (Previously Presented) The method according to claim 66, further comprising the step of adding a suspension of cells to a surface of the at least partially gelled first layer, before the step of placing the volume of the second polymerizable mixture in the tissue space.

77. (Previously Presented) The method according to claim 66, wherein at least one of the first, second, and third polymerizable mixtures further comprises a bioactive substance selected from the group consisting of a nutrient, a cellular mediator, a growth factor, a compound which induces cellular differentiation, a bioactive polymer, a gene vector, and a pharmaceutical.

78 to 87. (Canceled)